

Electroweak Baryogenesis with Lepton Flavor Violation

Cheng-Wei Chiang^{1,2,3,4,*}, Kaori Fuyuto^{5,†} and Eibun Senaha^{1,2‡}

¹*Department of Physics, National Taiwan University, Taipei 10617, Taiwan*

²*Department of Physics and Center for Mathematics and Theoretical Physics, National Central University, Taoyuan 32001, Taiwan*

³*Institute of Physics, Academia Sinica, Taipei 11529, Taiwan*

⁴*Physics Division, National Center for Theoretical Sciences, Hsinchu 30013, Taiwan and*

⁵*Department of Physics, Saga University, Saga 840-8502, Japan*

(Dated: October 4, 2016)

We investigate the feasibility of electroweak baryogenesis in a two-Higgs doublet model with lepton flavor violation. By scrutinizing the heavy Higgs boson mass spectrum, regions satisfying both strong first-order electroweak phase transition and the muon $g - 2$ anomaly are identified. We also estimate the baryon number density by exploiting extra Yukawa couplings in the μ - τ sector. It is found that a CP-violating source term can be enhanced by the μ - τ flavor-violating coupling together with the extra τ coupling. With $\mathcal{O}(1)$ Yukawa couplings and CP-violating phases, the observed baryon number density is marginally produced under a generous assumption for the bubble wall profile.

I. INTRODUCTION

Baryon asymmetry of the Universe (BAU) is an observational fact [1] whose origin still remains open and calls for physics beyond the standard model (SM). Two important ingredients, namely, CP violation and departure from thermal equilibrium, are known to be insufficient in the SM [2, 3]. A great number of baryogenesis mechanisms have been proposed so far, and the relevant energy scales are highly model-dependent. From the view point of testability, electroweak baryogenesis (EWBG) [4, 5] is the most attractive scenario since it can be probed by current and foreseeable future experiments. It is thus interesting and important to scrutinize its feasible parameter space.

One of the simplest extensions for successful EWBG is to add another Higgs doublet to the SM Higgs sector, rendering the so-called two-Higgs doublet model (2HDM) (for a review, see Ref. [6] and references therein). In this model, both Higgs doublets can couple to quarks and leptons concurrently, giving rise to Higgs-mediated flavor-changing processes at tree level. Importantly, flavor-changing neutral Higgs (FCNH) couplings are in general complex and may yield CP violation relevant to baryogenesis. Earlier studies on EWBG with FCNH interactions can be found in Refs. [7–9]. Moreover, it is possible to have a strong first-order electroweak phase transition (EWPT) owing to the contributions of the additional Higgs doublet [10–13].

Recently, the CMS Collaboration reported an excess in a lepton flavor-violating (LFV) Higgs decay, $\text{Br}(h \rightarrow \mu\tau) = (0.84^{+0.39}_{-0.37})\%$, showing a 2.4σ deviation from the SM [14].¹ The ATLAS Collaboration, on the other hand, quotes an upper bound $\text{Br}(h \rightarrow \mu\tau) < 1.43\%$ [16]. The

authors of Refs. [17, 18], including one of the present authors, pointed out that the 2HDM with μ - τ flavor violation granted a parameter space to explain not only the $h \rightarrow \mu\tau$ excess but also the long-standing anomaly in muon $g - 2$ without having conflicts with other LFV constraints, such as $\tau \rightarrow \mu\gamma$, $\tau \rightarrow \mu\nu\bar{\nu}$, etc. Since such FCNH interactions are generically accompanied by additional CP-violating sources, it is an interesting question whether the new CP-violating phases play a crucial role in generating the BAU.

In this paper, we examine the possibility of EWBG in the 2HDM with μ - τ flavor violation. Our analysis is twofold: (1) baryon number generation via the new CP violation and (2) baryon number preservation by a strong first-order EWPT. For the former, we evaluate the CP-violating source based on the method of closed-time path formalism with vacuum expectation value (VEV) insertion, and then solve the diffusion equations to estimate the BAU. For the latter, we employ the one-loop finite temperature effective potential with a thermal resummation to determine the order of EWPT. To determine the strength of first-order EWPT, we explicitly evaluate the energy of sphaleron configuration.

The paper is organized as follows. In Sec. II, we give basic ingredients of the 2HDM and show a relationship between FCNH couplings in the symmetric phase and those in the broken phase. In Sec. III, we review how $h \rightarrow \mu\tau$ and $(g - 2)_\mu$ can be simultaneously explained by the μ - τ flavor violation, as found in Refs. [17, 18]. The baryon number preservation condition is given in Sec. IV. We identify the regions where both strong first-order EWPT and $(g - 2)_\mu$ anomaly can be accounted for. In Sec. V, we present the BAU calculation and its numerical analysis. Our conclusion is given in Sec. VI.

* chengwei@phys.ntu.edu.tw

† ss6206@cc.saga-u.ac.jp

‡ senaha@ncu.edu.tw

¹ The excess would be less significant if the latest LHC Run-II data

are taken into account [15]. Statistically, it is not yet sufficient to claim that the excess is completely gone.

II. MODEL

The 2HDM is an extension of the SM by adding one additional Higgs doublet. Such a two-Higgs doublets structure is motivated by some UV theories such as supersymmetric theories. The most general Higgs potential at the renormalizable level is

$$\begin{aligned}
V_0(\Phi_1, \Phi_2) &= m_1^2 \Phi_1^\dagger \Phi_1 + m_2^2 \Phi_2^\dagger \Phi_2 - (m_3^2 \Phi_1^\dagger \Phi_2 + \text{h.c.}) \\
&+ \frac{\lambda_1}{2} (\Phi_1^\dagger \Phi_1)^2 + \frac{\lambda_2}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) \\
&+ \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) + \left[\frac{\lambda_5}{2} (\Phi_1^\dagger \Phi_2)^2 \right. \\
&\left. + \left\{ \lambda_6 (\Phi_1^\dagger \Phi_1) + \lambda_7 (\Phi_2^\dagger \Phi_2) \right\} (\Phi_1^\dagger \Phi_2) + \text{h.c.} \right]. \quad (1)
\end{aligned}$$

Hermiticity of V_0 requires that $m_1^2, m_2^2, \lambda_1, \lambda_2, \lambda_3$ and λ_4 are all real while $m_3^2, \lambda_5, \lambda_6$ and λ_7 are generally complex, with one of them being possibly made real by a field redefinition of either Higgs doublet. We parametrize Φ_i ($i = 1, 2$) as

$$\Phi_i(x) = e^{i\theta_i} \begin{pmatrix} \phi_i^+(x) \\ \frac{1}{\sqrt{2}} (v_i + h_i(x) + ia_i(x)) \end{pmatrix}, \quad (2)$$

where $v_i e^{i\theta_i}$ are the VEV's. For simplicity, we assume that CP is not violated by the Higgs potential and the VEV's, leading to $\theta_i = 0$. Here we express $v_{1,2}$ in terms of polar coordinates, $v_1 = v \cos \beta$ and $v_2 = v \sin \beta$ with $v \simeq 246$ GeV. In the following, we will use the shorthand notation $s_\beta = \sin \beta$, $c_\beta = \cos \beta$ and $t_\beta = \tan \beta$.

In phenomenological discussions, it is useful to use the Higgs (Georgi) basis [19] in which only one Higgs doublet develops the VEV and the Nambu-Goldstone bosons ($G^0, G^\pm$) are decoupled from the physical states:

$$\begin{aligned}
\Phi'_1 &= c_\beta \Phi_1 + s_\beta \Phi_2 = \begin{pmatrix} G^+ \\ \frac{1}{\sqrt{2}} (v + h'_1 + iG^0) \end{pmatrix}, \\
\Phi'_2 &= -s_\beta \Phi_1 + c_\beta \Phi_2 = \begin{pmatrix} H^+ \\ \frac{1}{\sqrt{2}} (h'_2 + iA) \end{pmatrix}, \quad (3)
\end{aligned}$$

where $h'_1 = c_{\beta-\alpha} H + s_{\beta-\alpha} h$ and $h'_2 = -s_{\beta-\alpha} H + c_{\beta-\alpha} h$ with α being a mixing angle between the two CP-even Higgs fields ($h_{1,2}$). In this paper, h is the 125 GeV Higgs boson and assumed to be the lighter one.

Without imposing any symmetries, both Higgs doublets can couple to fermions. The relevant Yukawa inter-

actions in the lepton sector are

$$\begin{aligned}
-\mathcal{L}_Y &= \bar{l}_{iL} (Y_1 \Phi_1 + Y_2 \Phi_2)_{ij} e_{jR} + \text{h.c.} \\
&\ni \bar{e}_{iL} \left[\frac{y_i}{\sqrt{2}} \delta_{ij} s_{\beta-\alpha} + \frac{1}{\sqrt{2}} \rho_{ij} c_{\beta-\alpha} \right] e_{jR} h \\
&+ \bar{e}_{iL} \left[\frac{y_i}{\sqrt{2}} \delta_{ij} c_{\beta-\alpha} - \frac{1}{\sqrt{2}} \rho_{ij} s_{\beta-\alpha} \right] e_{jR} H \\
&+ \frac{i}{\sqrt{2}} \bar{e}_{iL} \rho_{ij} e_{jR} A + \text{h.c.}, \quad (4)
\end{aligned}$$

where i, j are flavor indices, $Y_{1,2}$ are general 3-by-3 complex matrices, and

$$\rho_{ij} = -t_\beta y_i \delta_{ij} + \frac{1}{c_\beta} \left(V_L^{e\dagger} Y_2 V_R^e \right)_{ij}, \quad (5)$$

with $V_{L,R}^e$ defined as the unitary matrices that diagonalize the charged lepton Yukawa couplings $Y^{\text{SM}} = (Y_1 c_\beta + Y_2 s_\beta)$, *i.e.*, $V_L^{e\dagger} Y^{\text{SM}} V_R^e = Y_D = \text{diag}(y_e, y_\mu, y_\tau)$. After this diagonalization, the mass terms of the charged leptons are given by $m_i = y_i v / \sqrt{2}$ with $i = e, \mu, \tau$.

Non-diagonal elements of ρ_{ij} are the sources of tree-level FCNH processes. In the literature, the so-called Cheng-Sher ansatz [20] for ρ_{ij} is adopted in order to avoid experimental constraints. In our analysis, however, we will not assume it while obtaining a parameter space that can accommodate both $h \rightarrow \mu\tau$ and $g - 2$ anomalies. In general, $Y_{1,2}$ cannot be uniquely determined even though Y_D is known. In our analysis, we make some working assumption on $Y_{1,2}$ for baryogenesis, and determine the structure of FCNH couplings ρ at $T = 0$. For later use, we define $\rho_{ij} = |\rho_{ij}| e^{i\phi_{ij}}$.

III. $h \rightarrow \mu\tau, (g - 2)_\mu$ AND EDM

In this section, we outline some important consequences of Ref. [17] to make this paper self-contained.

The Higgs decay to μ and τ in the current model occurs at tree level through μ - τ interactions (for earlier studies, see Refs. [21]), and its branching ratio is given by

$$\text{Br}(h \rightarrow \mu\tau) = \frac{m_h (|\rho_{\mu\tau}|^2 + |\rho_{\tau\mu}|^2) c_{\beta-\alpha}^2}{16\pi\Gamma_h}, \quad (6)$$

where $m_h = 125$ GeV and $\Gamma_h = 4.1$ MeV. It is easy to accommodate $\text{Br}(h \rightarrow \mu\tau) \simeq 0.84\%$ by taking $|\rho_{\mu\tau}| \sim |\rho_{\tau\mu}| \sim \mathcal{O}(0.1)$ and $c_{\beta-\alpha} \sim \mathcal{O}(0.01)$. It had been shown that such parameter choices did not violate the current experimental bounds on other LFV processes such as $\tau \rightarrow \mu\gamma$, $\tau \rightarrow \mu\nu\bar{\nu}$, etc [17]. Note that it is sufficient for either $\rho_{\mu\tau}$ or $\rho_{\tau\mu}$ to be nonzero to explain the $h \rightarrow \mu\tau$ excess only. If both couplings coexist, on the other hand, we can have a one-loop diagram that induces $g - 2$ and electric dipole moment (EDM) of the muon, denoted by δa_μ and d_μ , respectively. Contributions of the neutral

Higgs bosons to δa_μ and d_μ are given by

$$\delta a_\mu = \frac{m_\mu m_\tau \text{Re}(\rho_{\mu\tau} \rho_{\tau\mu})}{16\pi^2} \times \left[\frac{c_{\beta-\alpha}^2 f(r_h)}{m_h^2} + \frac{s_{\beta-\alpha}^2 f(r_H)}{m_H^2} - \frac{f(r_A)}{m_A^2} \right], \quad (7)$$

$$\frac{d_\mu}{|e|} = -\frac{1}{2m_\mu} \text{Arg}(\rho_{\mu\tau} \rho_{\tau\mu}) \delta a_\mu, \quad (8)$$

where

$$f(r_\phi) = \frac{-1}{2(1-r_\phi)^2} \left[\frac{2 \ln r_\phi}{1-r_\phi} + 3 - r_\phi \right] \\ \underset{r_\phi \ll 1}{\simeq} \ln r_\phi^{-1} - \frac{3}{2}, \quad (9)$$

with $r_\phi = m_\tau^2/m_\phi^2$. Non-degeneracy in the neutral Higgs masses and a proper choice of the overall sign are essential for obtaining a sufficient δa_μ . In Ref. [22], the discrepancy of $(g-2)_\mu$ between experiment and the SM prediction was estimated as

$$\delta a_\mu = a_\mu^{\text{EXP}} - a_\mu^{\text{SM}} = (26.1 \pm 8.0) \times 10^{-10}. \quad (10)$$

As demonstrated in Ref. [17], this deviation could be accommodated for $m_{H,A,H^\pm} \in [200, 500]$ GeV with an appropriate mass hierarchy. From the EWBG point of view, this mass range of the heavy Higgs bosons has the right scale for realizing the first-order EWPT, though the LFV interactions by themselves do not play a central role in the realization, as will be shown in the following sections.

The current limit on d_μ is [23]

$$|d_\mu| < 1.9 \times 10^{-19} \text{ e cm}, \quad (11)$$

which is about three orders of magnitude larger than d_μ estimated with $\text{Arg}(\rho_{\mu\tau} \rho_{\tau\mu}) = 1$ and $\delta a_\mu = 3.0 \times 10^{-9}$ in Eq. (8). In what follows, we will clarify the relationship between CP violation appearing in d_μ and that relevant to BAU.

IV. BARYON NUMBER PRESERVATION

The baryon number is generated via the sphaleron process outside the bubble (symmetric phase), and it can survive if the sphaleron process is sufficiently quenched inside the bubble (broken phase). To this end, the sphaleron rate in the broken phase, denoted as $\Gamma_B^{(b)}(T)$, must be smaller than the Hubble constant, $H(T)$. More explicitly,

$$\Gamma_B^{(b)}(T) \simeq (\text{prefactor}) e^{-E_{\text{sph}}(T)/T} \\ < H(T) \simeq 1.66 \sqrt{g_*(T)} T^2 / m_{\text{P}}, \quad (12)$$

where E_{sph} stands for the sphaleron energy, g_* is the number of relativistic degrees of freedom in the plasma ($g_* = 110.75$ in the 2HDM), and $m_{\text{P}} = 1.22 \times 10^{19}$ GeV.

We parametrize the sphaleron energy as $E_{\text{sph}}(T) = 4\pi v(T) \mathcal{E}(T) / g_2$ with g_2 being the $\text{SU}(2)_L$ gauge coupling constant. Eq. (12) is then rewritten as

$$\frac{v(T)}{T} > \frac{g_2}{4\pi \mathcal{E}(T)} [42.97 + \log \text{ terms}] \equiv \zeta_{\text{sph}}(T). \quad (13)$$

One can see that $\zeta_{\text{sph}}(T)$ is mostly controlled by $\mathcal{E}(T)$. The logarithmic corrections in the bracket come from the prefactor of $\Gamma_B^{(b)}$. To our best knowledge, an extensive study on the prefactor in the 2HDM is still missing. In the minimal supersymmetric SM case, on the other hand, the zero mode factors of the fluctuations about the sphaleron typically amount to about 10% [24]. This is subdominant and, therefore, we will neglect them in our numerical analysis for simplicity.

We impose Eq. (13) at a critical temperature T_C at which the effective potential has two degenerate minima. In our analysis, T_C and $v_C \equiv v(T_C)$ are determined using finite-temperature one-loop effective potential with thermal resummation. As mentioned in Sec. III, $s_{\beta-\alpha}$ is close to 1 in the region of interest to us. In such a case, we may simplify the analysis of EWPT to a one-dimensional problem with a single order parameter, as discussed in Refs. [10, 12].

As is well-known, the extra heavy Higgs bosons can play a major role in enhancing v_C/T_C in the 2HDM. In this case, $M^2 \equiv m_3^2/(s_\beta c_\beta)$ must not exceed certain values, depending on the magnitude of quartic couplings; otherwise, the so-called non-decoupling effects would diminish, rendering a suppressed v_C/T_C . Phenomenological consequences of the non-decoupling effects at $T = 0$ include significant deviations in the $h \rightarrow \gamma\gamma$ decay width [25] and the triple Higgs coupling [26].

For the evaluation of $\mathcal{E}$, we solve the equations of motion for the sphaleron with appropriate boundary conditions [27, 28]. Here, we use the tree-level Higgs potential for simplicity. In this case, ζ_{sph} may be underestimated by $\mathcal{O}(10)\%$ since $\mathcal{E}(0) > \mathcal{E}(T_C)$. As will be shown below, this approximation does not affect our conclusion. A detailed analysis of $\zeta_{\text{sph}}(T)$ will be given elsewhere.

We note in passing that recent studies show that $\zeta_{\text{sph}}(T_C) = 1.1 - 1.2$ in the real singlet-extended SM [29] and 1.23 in the scale-invariant 2HDM [13] for typical parameter sets.

In Fig. 1, v_C/T_C and δa_μ are shown in the (m_H, m_A) plane. We take $m_A = m_{H^\pm}$ to avoid the electroweak ρ parameter constraint, and choose $c_{\beta-\alpha} = 0.006$, $|\rho_{\tau\mu}| = |\rho_{\mu\tau}|$, and $\phi_{\tau\mu} + \phi_{\mu\tau} = \pi/4$, as favored by the solution of $(g-2)_\mu$ anomaly. For the remaining parameters, we set $M = 100$ GeV, $\tan\beta = 1$ and $\lambda_6 = \lambda_7 = 0$ as an example. Contours of v_C/T_C are drawn with the solid gray curves with values of 1.0, 1.17, 1.5, 2.0, 2.5, and 3.0 from bottom to top, where 1.17 corresponds to $\zeta_{\text{sph}}(T_C)$. Allowed 1σ , 2σ and 3σ regions of δa_μ are shown by the areas colored in green, blue and pink, respectively. One can see that the regions satisfying the baryon number preservation condition ($v_C/T_C > \zeta_{\text{sph}}$) and favored by $(g-2)_\mu$ have an overlap if $m_A > m_H$. We note that




FIG. 1. Contours of v_C/T_C and δa_μ on the m_H - m_A plane. We take $m_A = m_{H^\pm}$, $M = 100$ GeV, $\tan \beta = 1$, $c_{\beta-\alpha} = 0.006$, $|\rho_{\tau\mu}| = |\rho_{\mu\tau}|$, and $\phi_{\tau\mu} + \phi_{\mu\tau} = \pi/4$. The solid gray curves are contours of $v_C/T_C = 1.0, 1.17 (= \zeta_{\text{sph}}(T_C)), 1.5, 2.0, 2.5$, and 3.0 from bottom to top. Areas colored in green, blue and pink correspond to the $1\sigma, 2\sigma$ and 3σ regions of δa_μ , respectively.

the $(g-2)_\mu$ -favored parameter space can change to the $m_A < m_H$ region if $\text{Re}(\rho_{\tau\mu}\rho_{\mu\tau}) < 0$, as inferred from Eq. (7). However, the allowed region shrinks in this case.

V. BARYON NUMBER GENERATION

We closely follow the method given in Refs. [30–33] (see also Refs. [34–36]) in estimating the BAU. In our scenario, the CP-violating source term is induced by interactions between chiral fermions and spacetime-dependent VEV's $v_i(X)$. As found in Ref. [37], the treatment of the chiral fermion should be taken with some care since its dispersion relation can be significantly modified by thermal plasma. As a result, the CPV source term for the left-handed (LH) fermion of flavor i induced by the right-handed (RH) fermion of flavor j takes the form ²

$$S_{iLjR}(X) = 2C(X) \int_0^\infty \frac{dk}{2\pi^2} k^2 \text{Im} \left[\frac{(1 - n_{jR}^p - n_{iL}^p) Z_{jR}^p Z_{iL}^p}{(\mathcal{E}_{jR}^p + \mathcal{E}_{iL}^p)^2} + \frac{n_{iL}^{h*} Z_{jR}^p Z_{iL}^{h*}}{(\mathcal{E}_{jR}^p - \mathcal{E}_{iL}^{h*})^2} + \frac{n_{jR}^{p*} Z_{jR}^{p*} Z_{iL}^h}{(\mathcal{E}_{jR}^{p*} - \mathcal{E}_{iL}^h)^2} + (p \leftrightarrow h) \right], \quad (14)$$

where $C(X) = |Y_{1ij}| |Y_{2ij}| \sin \theta_{ij} v^2(X) \partial_{t_X} \beta(X)$ with $\theta_{ij} = \text{Arg}(Y_{1ij}) - \text{Arg}(Y_{2ij})$, p denotes the ‘particle’ mode and h the ‘hole’ mode whose dispersion relations are

given by $\mathcal{E}^{p,h}(k) = E^{p,h}(k) - i\Gamma^{p,h}(k)$, and $Z_{iL,R}^p$ denote the residues. ³

Here, we remark on the treatment of $\Gamma^{p,h}$. In our numerical calculation, $\Gamma^{p,h}$ are approximated as $(\text{constant}) \times T$, as is often done in the calculation of the source terms for scalars and Dirac fermions. In the chiral fermion case, however, the term without statistical factor in the first term of Eq. (14) could cause a logarithmic divergence if Γ^p is fixed to a constant in the whole range of the k integration, which invalidates the calculation. Since the correct behavior of Γ^p in the large k region may take the form of $T^3 \ln(k/T)/k^2$ [40], the integration would be convergent. As a simple prescription, we take such a k -dependent Γ^h for large k . Ambiguities coming from this prescription will be discussed below.

One can also find that $S_{jRiL} = -S_{iLjR}$. In our study, we consider the case where the 32 and 33 elements of $Y_{1,2}$ play a dominant role in generating the BAU. In this case, only $S_{\tau_L\mu_R} (= -S_{\mu_R\tau_L})$ is relevant. As mentioned in Sec. II, $Y_{1,2}$ are connected with the $T = 0$ observables through Y_D and Eq. (5). For illustration, we assume that the 12, 13, 21 and 31 elements of Y^{SM} are all zero. Also, we focus on the case where only $\rho_{\mu\tau}$, $\rho_{\tau\mu}$ and $\rho_{\tau\tau}$ take nonzero values among ρ_{ij} .

Let us denote the relevant particle number densities as $Q_3 = n_{t_L} + n_{b_L}$, $T = n_{t_R}$, $B = n_{b_R}$, $L_3 = n_{\nu_{\tau L}} + n_{\tau_L}$, $R_3 = n_{\tau_R}$, $R_2 = n_{\mu_R}$, and $H = n_{H^+} + n_{H_1^0} + n_{H_2^+} + n_{H_2^0}$. The number density expanded to the leading order in the chemical potential μ is reduced to $n_{b,f} = T^2 \mu k_{b,f}/6$, with b (f) denoting bosons (fermions). In the massless case, one finds $k_b = 2$ and $k_f = 1$. For massive cases, more precise form of $k_{b,f}$ [32] should be used.

The set of Boltzmann equations may be cast into:

$$\begin{aligned} \partial_\mu j_{Q_3}^\mu &= -\Gamma_{Y_t}(\xi_{Q_3} + \xi_H - \xi_T) + \Gamma_{M_t}(\xi_T - \xi_{Q_3}) - 2\Gamma_{ss}N_5, \\ \partial_\mu j_T^\mu &= \Gamma_{Y_t}(\xi_{Q_3} + \xi_H - \xi_T) - \Gamma_{M_t}(\xi_T - \xi_{Q_3}) + \Gamma_{ss}N_5, \\ \partial_\mu j_H^\mu &= -\Gamma_{Y_t}(\xi_{Q_3} + \xi_H - \xi_T) + \Gamma_{Y_\tau}(\xi_{L_3} - \xi_H - \xi_{R_3}) \\ &\quad + \Gamma_{Y_{\tau\mu}}(\xi_{L_3} - \xi_H - \xi_{R_2}) - \Gamma_H \xi_H, \\ \partial_\mu j_{L_3}^\mu &= -\Gamma_{Y_\tau}(\xi_{L_3} - \xi_H - \xi_{R_3}) + \Gamma_{M_\tau}(\xi_{R_3} - \xi_{L_3}) \\ &\quad + \Gamma_{M_{\tau\mu}}^+(\xi_{R_2} + \xi_{L_3}) + \Gamma_{M_{\tau\mu}}^-(\xi_{R_2} - \xi_{L_3}) \\ &\quad - \Gamma_{Y_{\tau\mu}}(\xi_{L_3} - \xi_H - \xi_{R_2}) + S_{\tau_L\tau_R} + S_{\tau_L\mu_R}, \\ \partial_\mu j_{R_3}^\mu &= \Gamma_{Y_\tau}(\xi_{L_3} - \xi_H - \xi_{R_3}) - \Gamma_{M_\tau}(\xi_{R_3} - \xi_{L_3}) - S_{\tau_L\tau_R}, \\ \partial_\mu j_{R_2}^\mu &= \Gamma_{Y_{\tau\mu}}(\xi_{L_3} - \xi_H - \xi_{R_2}) \\ &\quad - \Gamma_{M_{\tau\mu}}^+(\xi_{R_2} + \xi_{L_3}) - \Gamma_{M_{\tau\mu}}^-(\xi_{R_2} - \xi_{L_3}) - S_{\tau_L\mu_R}, \end{aligned} \quad (15)$$

where $\xi_i = n_i/k_i$, $N_5 = 2\xi_{Q_3} - \xi_T + 9(Q_3 + T)/k_B$, and $\partial_\mu j_i^\mu = \dot{n}_i - D_i \nabla^2 n_i$ with D_i being a diffusion constant. In solving these coupled equations, we utilize the chemical equilibrium conditions in light of $\Gamma_{Y_t}^{-1}, \Gamma_{Y_\tau}^{-1}, \Gamma_{Y_{\tau\mu}}^{-1} <$

² Note that our result is different from the one shown in Ref. [38].

³ $\mathcal{E}^{p,h}(k)$ can be expressed in terms of the Lermbert W functions (for details, see Ref. [39]).

$T_C = 99.2 \text{ GeV}$	$v_C = 214.9 \text{ GeV}$	$v_w = 0.1$	$\Delta\beta = 0.05$
$m_{\tau_L} = 0.21T$	$m_{\mu_R} = 0.12T$	$\Gamma_{\tau_L} = 0.034T$	$\Gamma_{\mu_R} = 0.015T$
$D_q = 8.9/T$	$D_h = 101.9/T$	$D_L = 101.9/T$	$D_R = 377.1/T$

TABLE I. Input parameters for the BAU estimate.

τ_{diff} , the typical diffusion time for the particles under consideration. In this case, the above coupled Boltzmann equations are reduced to a single differential equation with respect to H , as is the case in Ref. [30].

The total LH number density (n_L) is

$$\begin{aligned} n_L &= 5Q_3 + 4T + L_3 \\ &= -\frac{9k_{Q_3}k_T - 5k_{Q_3}k_B - 8k_Tk_B}{k_H(9k_{Q_3} + 9k_T + k_B)} \\ &\quad + \frac{k_{L_3}D_R(k_{R_3} + k_{R_2})}{k_H(D_Lk_{L_3} + D_R(k_{R_3} + k_{R_2}))}, \end{aligned} \quad (16)$$

where D_L and D_R denote the diffusion constants of the third generation LH lepton doublet and the RH muon, respectively. As pointed out in Ref. [33], the lepton transport is much more efficient since it does not suffer from the strong sphaleron washout effect, which would make the first term highly suppressed.

With the above n_L , the BAU can be estimated as

$$n_B = \frac{-3\Gamma_B^{(s)}}{2D_q\lambda_+} \int_{-\infty}^0 dz' n_L(z') e^{-\lambda_- z'}, \quad (17)$$

where $\lambda_{\pm} = [v_w \pm \sqrt{v_w^2 + 4\mathcal{R}D_q}]/2D_q$, v_w represents the expanding speed of the bubble wall, D_q is the diffusion constant of the quarks, $\Gamma_B^{(s)}$ is the sphaleron rate in the unbroken phase, and $\mathcal{R} = 15\Gamma_B^{(s)}/4$. In the following, we will characterize the baryon asymmetry by $Y_B \equiv n_B/s$, where $s = (2\pi^2/45)g_*$ is the entropy density at temperature T , and quote $Y_B^{\text{obs}} = 8.59 \times 10^{-11}$, the central value given by the Planck Collaboration [1].

In the estimate of BAU, we employ the formulas given in Ref. [41] for the diffusion constants and thermal widths of LH and RH leptons, and assume $D_h = D_L$. The particular values that we use here are summarized in Table. I.

In Fig. 2, we show Y_B/Y_B^{obs} , $\text{Br}(h \rightarrow \mu\tau)$ and δa_μ as functions of $|\rho_{\tau\tau}|$ and $|\rho_{\tau\mu}|$. Here we take the same input parameters as those in Fig. 1 and the heavy Higgs boson masses $m_H = 350 \text{ GeV}$ and $m_A = m_{H^\pm} = 400 \text{ GeV}$, leading to $v_C/T_C = 214.9 \text{ GeV}/99.2 \text{ GeV} = 2.17$. As an ansatz for Y^{SM} , we consider

$$Y^{\text{SM}} = \begin{pmatrix} \sqrt{2}m_e/v & 0 & 0 \\ 0 & 3.31 \times 10^{-3} & -6.81i \times 10^{-4} \\ 0 & 8.91i \times 10^{-3} & 3.70 \times 10^{-3} \end{pmatrix}, \quad (18)$$




FIG. 2. Contours of Y_B/Y_B^{obs} , $\text{Br}(h \rightarrow \mu\tau)$ and δa_μ in the $(|\rho_{\tau\tau}|, |\rho_{\tau\mu}|)$ plane. We set $m_H = 350 \text{ GeV}$, $m_A = m_{H^\pm} = 400 \text{ GeV}$, $M = 100 \text{ GeV}$, $c_{\beta-\alpha} = 0.006$, $|\rho_{\tau\mu}| = |\rho_{\mu\tau}|$, $\phi_{\tau\mu} = -5\pi/4$, $\phi_{\mu\tau} = \pi/4 - \phi_{\tau\mu}$ and $\phi_{\tau\tau} = \pi/2$.

which is diagonalized by

$$V_R^e = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -0.365i & -0.931i \\ 0 & -0.931 & 0.365 \end{pmatrix}, \quad (19)$$

$$V_L^e = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -0.945i & -0.327i \\ 0 & -0.327 & 0.945 \end{pmatrix}. \quad (20)$$

Also, we take $\phi_{\tau\tau} = \pi/2$. Note that each of $Y_{1,2}$ is fixed by $V_L^{e\dagger} Y^{\text{SM}} V_R^e = Y_D$ and Eq. (5) once ρ_{ij} are given.

The solid lines represent the contours of $Y_B/Y_B^{\text{obs}} = 0.5$ (left) and 1.0 (right). The dashed lines in black and red represent $\text{Br}(h \rightarrow \mu\tau) = 1.43\%$ and 0.84% , respectively. The colored regions with the same color scheme as in Fig. 1 explain the $(g-2)_\mu$ anomaly. In our setup, the dominant contribution to the CP-violating source term comes from $S_{\tau_L\mu_R}$ which is induced by the 32 elements of Y_1 and Y_2 . We find that these off-diagonal elements have stronger dependences on $|\rho_{\tau\tau}|$ than $|\rho_{\tau\mu}|$ and $|\rho_{\mu\tau}|$. Under rather generous assumptions for the bubble wall profile, the generated BAU can reach its observed value for $|\rho_{\tau\mu}| \simeq 0.1 - 0.6$ and $|\rho_{\tau\tau}| \simeq 0.8 - 0.9$. Our main conclusion is that there is a parameter space that is consistent with both experimental anomalies of $h \rightarrow \mu\tau$ and $(g-2)_\mu$ as well as the observed BAU.

Some comments on the theoretical uncertainties are in order. (1) We take the same size of $|\Delta\beta|$ as in Ref. [9] as a reference value. However, quantitative studies of it in the 2HDM are still absent. In the MSSM, $\Delta\beta = \mathcal{O}(10^{-2} - 10^{-4})$ depending on m_A [42]. In the next-to-MSSM, $\Delta\beta$ can reach $\mathcal{O}(0.1)$ in some parameter space [43]. We should note that Y_B is mostly subject to the uncertainties of $\Delta\beta$ among others since it is linearly proportional to $\Delta\beta$. (2) Studies on v_w can be found in




FIG. 3. Correlations between BAU and electron EDM in the $(|\rho_{\tau\tau}|, \phi_{\tau\tau})$ plane. The contours shown in white are $|d_e| = 2.0 \times 10^{-29} e \text{ cm}$ (left) and $3.0 \times 10^{-29} e \text{ cm}$ (right). On the other hand, $Y_B/Y_B^{\text{obs}} = 0.5$ (left) and 1.0 (right) are denoted by the contours in black.

Refs. [44–47], which suggest that $0.1 \lesssim v_w \lesssim 0.6$ in non-SUSY models. For stronger first-order EWPT, v_w tends to be large and may reduce Y_B to some extent. In our analysis, we simply adopt the lowest value as the most generous choice. A more precise determination of v_w using our input parameters is definitely indispensable to obtaining more precise Y_B . (3) As mentioned above, the treatment of Γ^h is somewhat tricky. It is found that Y_B may change by a factor of a few or more, depending on at which scale the k -dependent Γ^h is put in. (4) The VEV-insertion method used here is vulnerable to theoretical uncertainties (see, *e.g.*, Refs. [5, 48]) and may lead to an overestimated BAU compared to an all-order VEV resummation method [49–51]. However, a satisfactory formalism of the Y_B calculation beyond that is still not available, and hence the error associated with the approximation cannot be properly quantified.

In summary, the observed Y_B can be marginally produced with the generous choice of the input parameters. However, a definitive statement cannot be made until the above-mentioned various theoretical uncertainties are fully under control.

We now turn to another experimental constraint. As $|\rho_{\tau\tau}|$ increases, $\text{Br}(\tau \rightarrow \mu\gamma)$ gets enhanced and, for $|\rho_{\tau\tau}| \gtrsim 0.1$, exceeds the current experimental upper bound, $\text{Br}(\tau \rightarrow \mu\gamma) < 4.4 \times 10^{-8}$ [52]. However, as demonstrated in Ref. [17], an accidental cancellation between the one-loop and two-loop contributions could occur in $\text{Br}(\tau \rightarrow \mu\gamma)$ if the new top Yukawa coupling (denoted as ρ_{tt}) took a nonzero value. In the current case, $|\rho_{tt}| \simeq 0.5$ with $\phi_{tt} \simeq \phi_{\tau\tau}$ gives $\text{Br}(\tau \rightarrow \mu\gamma) \simeq 2 \times 10^{-8}$ for $|\rho_{\tau\tau}| \simeq 1$. The impact of ϕ_{tt} on Y_B highly depends on the Yukawa structure of the top quark. With our as-

sumption, the top quark does not provide dominant CP violation for the baryogenesis.

Since $\rho_{\tau\tau}$ and ρ_{tt} are complex, they may induce an electron EDM through two-loop Barr-Zee diagrams, among which one Higgs-photon mediated loop diagram is dominant. In Fig. 3, $|d_e|$ and Y_B/Y_B^{obs} are plotted as functions of $|\rho_{\tau\tau}|$ and $\phi_{\tau\tau}$. The contours in white represent $|d_e| = 2.0 \times 10^{-29} e \text{ cm}$ (left) and $3.0 \times 10^{-29} e \text{ cm}$ (right). $|d_e|$ with $|\rho_{\tau\tau}| = 1$ and $\phi_{\tau\tau} = \pi/2$ reaches a maximal value of $3.5 \times 10^{-29} e \text{ cm}$ which is slightly smaller than the current bound on electron EDM, $|d_e| < 8.7 \times 10^{-29} e \text{ cm}$ [53]. This is due to the facts that the heavy Higgs boson couplings to the electron is suppressed by $c_{\beta-\alpha}$ and that the extra Yukawa coupling ρ_{ee} is absent. It should be noted that the complex phases of $\rho_{\tau\mu}$ and $\rho_{\mu\tau}$ do not contribute to d_e via the Higgs-photon mediated Barr-Zee diagrams since the internal photon line cannot change the fermion flavors. As discussed in Sec. III, on the other hand, the muon EDM is induced by those FCNH couplings at one-loop level, giving rise to $|d_\mu| \simeq 3 \times 10^{-22} e \text{ cm}$ for the current parameter set.

We also find that $\tau \rightarrow \mu\nu\bar{\nu}$ can give some constraints on $|\rho_{\tau\mu}|$ and $|\rho_{\mu\tau}|$ as well as $m_{H^\pm}$ and can be similar to the constraint coming from $\text{Br}(h \rightarrow \tau\mu) < 1.43$.

VI. CONCLUSION

We have studied electroweak baryogenesis in the general framework of the two-Higgs doublet model in light of the $h \rightarrow \mu\tau$ and $(g-2)_\mu$ anomalies. In this model, the heavy Higgs bosons with the appropriate μ - τ flavor violation can accommodate the above two anomalies. At the same time, these extra Higgs bosons can induce a strong first-order electroweak phase transition as required for successful electroweak baryogenesis.

It is found that the μ - τ flavor-violating lepton sector has a great potential to generate sufficient baryon asymmetry of the Universe within the theoretical uncertainties. In this scenario, the interplay between $\rho_{\tau\mu}/\rho_{\mu\tau}$ and $\rho_{\tau\tau}$ is crucially important. Our analysis suggests that $Y_B/Y_B^{\text{obs}} \simeq 1$ for $|\rho_{\tau\mu}| \simeq 0.1 - 0.6$ and $|\rho_{\tau\tau}| \simeq 0.8 - 0.9$ with $\mathcal{O}(1)$ CP-violating phases. To suppress $\text{Br}(\tau \rightarrow \mu\gamma)$, a cancellation mechanism has to be at work, which additionally predicts $|\rho_{tt}| \simeq 0.5$ and $\phi_{tt} \simeq \phi_{\tau\tau}$. Since future experimental sensitivities of $\text{Br}(\tau \rightarrow \mu\gamma)$ and $|d_e|$ are about 1×10^{-9} [54] and $10^{-30} e \text{ cm}$ [55], respectively, our scenario could be fully tested.

ACKNOWLEDGMENTS

E.S. thanks Yoshimasa Hidaka for useful discussions. This work is supported in part by the Ministry of Science and Technology of R.O.C. under Grant Nos. 104-2628-M-002-014-MY4 (CWC) and 104-2811-M-008-011 (ES), and in part by the Research Fellowships of the Japan Society

-
- [1] P. A. R. Ade *et al.* [Planck Collaboration], *Astron. Astrophys.* **571**, A16 (2014).
- [2] M. B. Gavela, P. Hernandez, J. Orloff and O. Pene, *Mod. Phys. Lett. A* **9**, 795 (1994); M. B. Gavela, P. Hernandez, J. Orloff, O. Pene and C. Quimbay, *Nucl. Phys. B* **430**, 382 (1994); P. Huet and E. Sather, *Phys. Rev. D* **51**, 379 (1995); T. Konstandin, T. Prokopec and M. G. Schmidt, *Nucl. Phys. B* **679**, 246 (2004).
- [3] K. Kajantie, M. Laine, K. Rummukainen and M. E. Shaposhnikov, *Phys. Rev. Lett.* **77**, 2887 (1996); K. Rummukainen, M. Tsy-pin, K. Kajantie, M. Laine and M. E. Shaposhnikov, *Nucl. Phys. B* **532**, 283 (1998); F. Csikor, Z. Fodor and J. Heitger, *Phys. Rev. Lett.* **82**, 21 (1999); Y. Aoki, F. Csikor, Z. Fodor and A. Ukawa, *Phys. Rev. D* **60**, 013001 (1999).
- [4] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, *Phys. Lett. B* **155** (1985) 36. For reviews on electroweak baryogenesis, see A. G. Cohen, D. B. Kaplan and A. E. Nelson, *Ann. Rev. Nucl. Part. Sci.* **43** (1993) 27; M. Quiros, *Helv. Phys. Acta* **67** (1994) 451; V. A. Rubakov and M. E. Shaposhnikov, *Usp. Fiz. Nauk* **166** (1996) 493; K. Funakubo, *Prog. Theor. Phys.* **96** (1996) 475; M. Trodden, *Rev. Mod. Phys.* **71** (1999) 1463; W. Bernreuther, *Lect. Notes Phys.* **591** (2002) 237; J. M. Cline, [arXiv:hep-ph/0609145]; T. Konstandin, *Phys. Usp.* **56** (2013) 747 [*Usp. Fiz. Nauk* **183** (2013) 785].
- [5] D. E. Morrissey and M. J. Ramsey-Musolf, *New J. Phys.* **14**, 125003 (2012).
- [6] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher and J. P. Silva, *Phys. Rept.* **516** (2012) 1.
- [7] S. Tulin and P. Winslow, *Phys. Rev. D* **84** (2011) 034013.
- [8] J. M. Cline, K. Kainulainen and M. Trott, *JHEP* **1111** (2011) 089.
- [9] T. Liu, M. J. Ramsey-Musolf and J. Shu, *Phys. Rev. Lett.* **108** (2012) 221301.
- [10] A. I. Bochkarev, S. V. Kuzmin and M. E. Shaposhnikov, *Phys. Lett. B* **244** (1990) 275; N. Turok and J. Zadrozny, *Nucl. Phys. B* **369** (1992) 729; J. M. Cline and P. A. Lemieux, *Phys. Rev. D* **55** (1997) 3873.
- [11] D. Land and E. D. Carlson, *Phys. Lett. B* **292**, 107 (1992); A. Hammerschmitt, J. Kripfganz and M. G. Schmidt, *Z. Phys. C* **64**, 105 (1994); K. Funakubo, A. Kakuto and K. Takenaga, *Prog. Theor. Phys.* **91**, 341 (1994); L. Fromme, S. J. Huber and M. Seniuch, *JHEP* **0611** (2006) 038; G. Gil, P. Chankowski and M. Krawczyk, *Phys. Lett. B* **717** (2012) 396; J. M. Cline and K. Kainulainen, *Phys. Rev. D* **87** (2013) no.7, 071701; G. C. Dorsch, S. J. Huber and J. M. No, *JHEP* **1310** (2013) 029; G. C. Dorsch, S. J. Huber, K. Mimasu and J. M. No, *Phys. Rev. Lett.* **113** (2014) no.21, 211802; N. Blinov, S. Profumo and T. Stefaniak, *JCAP* **1507** (2015) no.07, 028;
- [12] S. Kanemura, Y. Okada and E. Senaha, *Phys. Lett. B* **606** (2005) 361.
- [13] K. Fuyuto and E. Senaha, *Phys. Lett. B* **747** (2015) 152.
- [14] V. Khachatryan *et al.* [CMS Collaboration], *Phys. Lett. B* **749**, 337 (2015)
- [15] Talk given by Maria Cepeda at Higgs Tasting 2016, Benasque, Spain.
- [16] G. Aad *et al.* [ATLAS Collaboration], arXiv:1604.07730 [hep-ex].
- [17] Y. Omura, E. Senaha and K. Tobe, *JHEP* **1505**, 028 (2015); Y. Omura, E. Senaha and K. Tobe, arXiv:1511.08880 [hep-ph].
- [18] K. Tobe, arXiv:1607.04447 [hep-ph].
- [19] H. Georgi and D. V. Nanopoulos, *Phys. Lett. B* **82**, 95 (1979); J. F. Donoghue and L. F. Li, *Phys. Rev. D* **19**, 945 (1979); L. Lavoura and J. P. Silva, *Phys. Rev. D* **50**, 4619 (1994); L. Lavoura, *Phys. Rev. D* **50**, 7089 (1994); F. J. Botella and J. P. Silva, *Phys. Rev. D* **51**, 3870 (1995); G. C. Branco, L. Lavoura and J. P. Silva, *Int. Ser. Monogr. Phys.* **103**, 1 (1999); S. Davidson and H. E. Haber, *Phys. Rev. D* **72**, 035004 (2005) Erratum: [*Phys. Rev. D* **72**, 099902 (2005)].
- [20] T. P. Cheng and M. Sher, *Phys. Rev. D* **35** (1987) 3484. doi:10.1103/PhysRevD.35.3484
- [21] A. Pilaftsis, *Phys. Lett. B* **285**, 68 (1992); J. G. Korner, A. Pilaftsis and K. Schilcher, *Phys. Rev. D* **47**, 1080 (1993); K. A. Assamagan, A. Deandrea and P. A. Delsart, *Phys. Rev. D* **67**, 035001 (2003); A. Brignole and A. Rossi, *Phys. Lett. B* **566**, 217 (2003); S. Kanemura, K. Matsuda, T. Ota, T. Shindou, E. Takasugi and K. Tsumura, *Phys. Lett. B* **599**, 83 (2004); E. Arganda, A. M. Curiel, M. J. Herrero and D. Temes, *Phys. Rev. D* **71**, 035011 (2005); S. Kanemura, T. Ota and K. Tsumura, *Phys. Rev. D* **73**, 016006 (2006); G. Blankenburg, J. Ellis and G. Isidori, *Phys. Lett. B* **712**, 386 (2014); R. Harnik, J. Kopp and J. Zupan, *JHEP* **1303**, 026 (2013); M. Arana-Catania, E. Arganda and M. J. Herrero, *JHEP* **1309**, 160 (2013) [*JHEP* **1510**, 192 (2015)]; E. Arganda, M. J. Herrero, X. Marcano and C. Weiland, *Phys. Rev. D* **91**, no. 1, 015001 (2015); E. Arganda, M. J. Herrero, X. Marcano and C. Weiland, *Phys. Rev. D* **93** (2016) no.5, 055010; E. Arganda, M. J. Herrero, R. Morales and A. Szykman, *JHEP* **1603** (2016) 055.
- [22] K. Hagiwara, R. Liao, A. D. Martin, D. Nomura and T. Teubner, *J. Phys. G* **38**, 085003 (2011).
- [23] G. W. Bennett *et al.* [Muon (g-2) Collaboration], *Phys. Rev. D* **80**, 052008 (2009).
- [24] K. Funakubo and E. Senaha, *Phys. Rev. D* **79** (2009) 115024.
- [25] I. F. Ginzburg, M. Krawczyk and P. Osland, In *Seogwipo 2002, Linear colliders* 90-94 [hep-ph/0211371].
- [26] S. Kanemura, S. Kiyoura, Y. Okada, E. Senaha and C. P. Yuan, *Phys. Lett. B* **558** (2003) 157; S. Kanemura, Y. Okada, E. Senaha and C.-P. Yuan, *Phys. Rev. D* **70** (2004) 115002.
- [27] N. S. Manton, *Phys. Rev. D* **28**, 2019 (1983).
- [28] F. R. Klinkhamer and N. S. Manton, *Phys. Rev. D* **30**, 2212 (1984).
- [29] K. Fuyuto and E. Senaha, *Phys. Rev. D* **90** (2014) no.1, 015015.
- [30] P. Huet and A. E. Nelson, *Phys. Rev. D* **53**, 4578 (1996).
- [31] A. Riotto, *Nucl. Phys. B* **518**, 339 (1998); A. Riotto,

- Phys. Rev. D **58**, 095009 (1998).
- [32] C. Lee, V. Cirigliano and M. J. Ramsey-Musolf, Phys. Rev. D **71**, 075010 (2005).
 - [33] D. J. H. Chung, B. Garbrecht, M. J. Ramsey-Musolf and S. Tulin, Phys. Rev. D **81** (2010) 063506
 - [34] K. Cheung, T. J. Hou, J. S. Lee and E. Senaha, Phys. Lett. B **710** (2012) 188.
 - [35] E. Senaha, Phys. Rev. D **88**, no. 5, 055014 (2013).
 - [36] K. Fuyuto, J. Hisano and E. Senaha, Phys. Lett. B **755**, 491 (2016).
 - [37] H. A. Weldon, Phys. Rev. D **61** (2000) 036003.
 - [38] K. Blum, C. Delaunay, M. Losada, Y. Nir and S. Tulin, JHEP **1005**, 101 (2010)
 - [39] C. P. Kiessig, M. Plumacher and M. H. Thoma, Phys. Rev. D **82** (2010) 036007.
 - [40] D. Boyanovsky, Phys. Rev. D **72** (2005) 033004.
 - [41] M. Joyce, T. Prokopec and N. Turok, Phys. Lett. B **338** (1994) 269; M. Joyce, T. Prokopec and N. Turok, Phys. Rev. D **53** (1996) 2930; M. Joyce, T. Prokopec and N. Turok, Phys. Rev. D **53** (1996) 2958.
 - [42] J. M. Moreno, M. Quiros and M. Seco, Nucl. Phys. B **526** (1998) 489.
 - [43] J. Kozaczuk, S. Profumo, L. S. Haskins and C. L. Wainwright, JHEP **1501** (2015) 144.
 - [44] A. Megevand and A. D. Sanchez, Nucl. Phys. B **825** (2010) 151.
 - [45] M. Ahmadvand, Int. J. Mod. Phys. A **29** (2014) no.20, 1450090
 - [46] T. Konstandin, G. Nardini and I. Rues, JCAP **1409** (2014) no.09, 028.
 - [47] J. Kozaczuk, JHEP **1510** (2015) 135.
 - [48] S. Inoue, G. Ovanessian and M. J. Ramsey-Musolf, Phys. Rev. D **93** (2016) 015013.
 - [49] M. Carena, J. M. Moreno, M. Quiros, M. Seco and C. E. M. Wagner, Nucl. Phys. B **599** (2001) 158.
 - [50] M. Carena, M. Quiros, M. Seco and C. E. M. Wagner, Nucl. Phys. B **650**, 24 (2003).
 - [51] T. Konstandin, T. Prokopec and M. G. Schmidt, Nucl. Phys. B **716**, 373 (2005); T. Konstandin, T. Prokopec, M. G. Schmidt and M. Seco, Nucl. Phys. B **738**, 1 (2006).
 - [52] K. Hayasaka *et al.* [Belle Collaboration], Phys. Lett. B **666** (2008) 16; B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. Lett. **104** (2010) 021802.
 - [53] J. Baron *et al.* [ACME Collaboration], Science **343**, 269 (2014) [arXiv:1310.7534 [physics.atom-ph]].
 - [54] T. Aushev *et al.*, arXiv:1002.5012 [hep-ex].
 - [55] Y. Sakemi, K. Harada, T. Hayamizu, M. Itoh, H. Kawamura, S. Liu, H. S. Nataraj and A. Oikawa *et al.*, J. Phys. Conf. Ser. **302** (2011) 012051; D. M. Kara, I. J. Smallman, J. J. Hudson, B. E. Sauer, M. R. Tarbutt and E. A. Hinds, New J. Phys. **14** (2012) 103051; D. Kawall, J. Phys. Conf. Ser. **295** (2011) 012031.